

## Estimating D.B.H. of Southern Indiana Oaks From Stump Diameter

Dale R. Weigel, *Forester* and Paul S. Johnson, *Principal Silviculturist*

### Application

Estimates of tree diameter at breast height (d.b.h.) based on measurements of diameter at stump height (d.s.h.) can be used to reconstruct preharvest volume, value, structure, and composition of stands after a timber harvest. It may be necessary or useful to do this in cases of timber theft and trespass or in certain silvicultural and ecological research applications. This report presents estimates of d.b.h. for five upland oak species in southern Indiana based on d.s.h. measured 6 inches above ground on the uphill side of the tree (table 1). The composite model (the average of all species) can be used when it is difficult or impossible to distinguish among species.

Other models (see references cited below) may be more appropriate for use outside of southern Indiana. Models applied outside their area of origin should be tested against a sample of at least 30 pairs of d.b.h./d.s.h. measurements that are well distributed across the d.b.h. range of each species of interest. If there is acceptable agreement between the model and the test data, application of the model is likely to produce acceptable estimates. Data on the agreement of the models presented herein with other published models are presented in Johnson and Weigel (1990).

### How The Models Were Developed

Because of the high correlation between d.b.h. and d.s.h., d.b.h. can be estimated with relatively high accuracy. We measured the d.b.h. and d.s.h. of 1,021 oaks of five species in 10 stands in southern Indiana. The number of trees measured per species ranged from 98 for northern red oak to 413 for chestnut oak. Measurements were made to the

nearest 0.04 inches at 4.5 feet above ground and at 6 inches above ground on the uphill side of each tree. Measurements were well distributed across the d.s.h. range of 3 to 36 inches in well-stocked upland hardwood stands; black oak site indices ranged from 60 to 75 feet at an index age of 50 years.

The models are given by the following non-linear regression equations:

•White oak-scarlet oak  
 $d.b.h. = 0.7586(d.s.h.)^{0.9900}$

•Black oak-chestnut oak  
 $d.b.h. = 0.9023(d.s.h.)^{0.9487}$

•Northern red oak  
 $d.b.h. = 0.9634(d.s.h.)^{0.9130}$

•Composite model  
 $d.b.h. = 0.8923(d.s.h.)^{0.9567}$

The  $R^2$ 's for all models were  $>0.96$ .

### References

- Bylin, Carl V. 1982. **Estimating dbh from stump diameter for 15 southern species.** Res. Note SO-286. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 3 p.
- Hampf, Frederick E. 1955. **Relationship of stump diameter to d.b.h. for northern red oak in the Northeast.** For. Res. Notes 46. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 3 p.

Table 1.—*D.b.h. of southern Indiana oaks estimated from stump diameter*

Stump diameter (in) <sup>1</sup>	White oak and scarlet oak	Black oak and chestnut oak	Northern red oak	Composite model
- - - - - <i>D.b.h. outside bark (inches)</i> - - - - -				
3	2.3	2.6	2.6	2.6
4	3.0	3.4	3.4	3.4
5	3.7	4.2	4.2	4.2
6	4.5	5.0	5.0	5.0
7	5.2	5.7	5.7	5.8
8	6.0	6.5	6.5	6.6
9	6.7	7.3	7.2	7.3
10	7.4	8.1	7.9	8.1
11	8.2	8.8	8.6	8.9
12	8.9	9.6	9.3	9.7
13	9.6	10.3	10.1	10.4
14	10.4	11.1	10.8	11.2
15	11.1	11.8	11.5	12.0
16	11.8	12.6	12.2	12.7
17	12.6	13.3	12.8	13.5
18	13.3	14.1	13.5	14.2
19	14.0	14.8	14.2	15.0
20	14.8	15.6	14.9	15.8
21	15.5	16.3	15.6	16.5
22	16.2	17.0	16.2	17.3
23	17.0	17.8	16.9	18.0
24	17.7	18.5	17.6	18.8
25	18.4	19.2	18.3	19.5
26	19.1	20.0	18.9	20.2
27	19.9	20.7	19.6	21.0
28	20.6	21.4	20.3	21.7
29	21.3	22.1	20.9	22.5
30	22.1	22.9	21.6	23.2
31	22.8	23.6	22.2	24.0
32	23.5	24.3	22.9	24.7
33	24.2	25.0	23.5	25.4
34	25.0	25.7	24.2	26.2
35	25.7	26.5	24.8	26.9
36	26.4	27.2	25.5	27.6

<sup>1</sup> Measured outside bark 6 inches above ground.

- Hampf, Frederick E. 1957. **Relationship of stump diameter to d.b.h. for white oak in the Northeast.** For. Res. Notes 66. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 3 p.
- Horn, Arthur G.; Keller, Richard C. 1957. **Tree diameter at breast height in relation to stump diameter by species group.** Tech. Notes 507. St. Paul, MN: U.S. Department of Agriculture, Forest Service, Lake States Forest Experiment Station. 2 p.
- Johnson, Paul S.; Weigel, Dale R. 1990. **Models for estimating DBH from stump diameter for southern Indiana oaks.** Northern Journal of Applied Forestry. 7(2): 79-81.
- McClure, Joe P. 1968. **Predicting tree D.B.H. from stump measurements in the Southeast.** Res. Note SE-99. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 4 p.
- Nyland, Ralph D. 1977. **DBH from stump diameter and height for northern hardwoods.** Res. Note 24. Syracuse, NY: State University of New York, College of Environmental Science and Forestry, Applied Forestry Research Institute. 2 p.
- Ojasvi, P.R.; Ramm, C.W.; Lantagne, D.O.; Bruggink, J. 1991. **Stump diameter and DBH relationships for white oak and black oak in the Lower Peninsula of Michigan.** Canadian Journal of Forest Research. 21: 1596-1600.
- Raile, Gerhard. 1978. **Estimating D.B.H. from stump dimensions.** In: Proceedings, 1977 Midwest forest mensurationists meeting; 1977; Lutsen, MN. Gen. Tech. Rep. NC-46. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station: 30-33.
- Vimmerstedt, J.P. 1957. **Estimating D.B.H. from stump diameter in southern Appalachian species.** Res. Notes 110. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 2 p.
- Wharton, Eric H. 1984. **Predicting diameter at breast height from stump diameters for northeastern tree species.** Res. Note NE-322. Broomall, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 4 p.
- Wiant, Harry V.; Williams, Thomas B. 1987. **Lower bole diameter and volume of four Appalachian hardwoods.** Northern Journal of Applied Forestry. 4(4): 212.

The United States Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs and marital or familial status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact the USDA Office of Communications at (202) 720-2791 (voice) or (202) 720-1127 (TDD).

To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call (800) 245-6340 (voice) or (202) 720-1127 (TDD). USDA is an equal employment opportunity employer.



North Central Forest Experiment Station  
1-26 Agriculture Building  
University of Missouri  
Columbia, Missouri 65211  
(573) 875-5341

